arXiv:hep-ph/9606451v1 27 Jun 1996

hep-ph/9606451

RECENT RESULTS IN THE NJL MODEL WITH HEAVY QUARKS*

THORSTEN FELDMANN†
Institute of Physics, Humboldt University, Invalidenstraße 110,
10115 Berlin, Germany

ABSTRACT

We investigate the interplay of chiral and heavy quark symmetries by using the NJL quark model. Heavy quarks with finite masses m_Q as well as the limit $m_Q \to \infty$ are studied. We found large corrections to the heavy mass scaling law for the pseudoscalar decay constant. The influence of external momenta on the shape parameters of the Isgur-Wise form factor is discussed.

1. Introduction

As is well known for heavy quark masses $m_Q \gg \Lambda_{QCD}$ one can expand the QCD lagrangian connected with heavy quark dynamics in terms of $1/m_Q$, leading to

$$\mathcal{L}_{\rm HQL} = \bar{Q}_v \, (iv\cdot D) \, Q_v + \mathcal{K}_v + \mathcal{M}_v + O(1/m_Q^2) \; , \tag{1}$$

with D_μ being the QCD covariant derivative and $\mathcal{K}_v$ and $\mathcal{M}_v$ being the kinetic and chromomagnetic energy of the heavy quark, respectively[1]. In the heavy quark limit $m_Q \to \infty$ (HQL) the contribution of the latter vanishes, and the remaining lagrangian is independent of the heavy quark flavor and spin. Consequently, heavy mesons are organized in spin symmetry doublets with $J = j_l \pm 1/2$ where j_l is the spin of the light degrees of freedom[2].

In the sector of light quark flavors $q = (u, d, s)$, QCD possesses an approximate $SU(3)_L \times SU(3)_R$ chiral symmetry which is spontaneously broken to $SU(3)_V$, leading to the emergence of (pseudo)Goldstone bosons π, K, η. We use the common non-linear representation, where the Goldstone bosons (denoted by π) and their transformations under chiral symmetry $SU(3)_L \times SU(3)_R$ are given by

$$\xi = e^{i\pi/F} \to L \, \xi \, U^\dagger = U \, \xi \, R^\dagger \; , \tag{2}$$

which defines the matrix $U(\pi, L, R)$ as a non-linear function of its arguments. Then heavy meson fields Φ transform under chiral symmetry as[3,4] $\Phi \to \Phi \, U^\dagger$.

---

[*] Talk presented at the III. German-Russian Workshop on *Theoretical Progress in* Heavy Quark Physics, Dubna, 20-22 May 1996. To appear in the conference proceedings.

[†] Supported by *Deutsche Forschungsgemeinschaft* under contract Eb 139/1–2.

2. The NJL model with heavy quarks

We will use the above symmetry considerations in order to construct the NJL quark model as a low-energy approximation to QCD. One choses suitable 4-quark operators representing the scalar, pseudoscalar, vector and axial-vector channel, respectively,

$$\begin{aligned} \mathcal{L}^{\bar{q}Q} &= 2\,G_3\,\Big((\bar{Q}\,q)\,(\bar{q}\,Q) + (\bar{Q}\,i\gamma_5\,q)\,(\bar{q}\,i\gamma_5\,Q) \\ &\qquad -\frac{1}{2}\,(\bar{Q}\,\gamma_\mu\,q)\,(\bar{q}\,\gamma^\mu\,Q) - \frac{1}{2}\,(\bar{Q}\,i\gamma_\mu\gamma_5\,q)\,(\bar{q}\,i\gamma_5\gamma^\mu\,Q)\Big) \\ \xrightarrow{m_Q\to\infty}\ \mathcal{L}^{\bar{q}Q_v} &= G_3\,\Big((\bar{Q}_v\,i\gamma_5\,q)\,(\bar{q}\,i\gamma_5\,Q_v) - (\bar{Q}_v\,\gamma^\perp_\mu\,q)\,(\bar{q}\,\gamma^\mu_\perp\,Q_v) \\ &\qquad +(\bar{Q}_v\,q)\,(\bar{q}\,Q_v) - (\bar{Q}_v\,i\gamma^\perp_\mu\gamma_5\,q)\,(\bar{q}\,i\gamma_5\gamma^\mu_\perp\,Q_v)\Big)\ . \end{aligned} \tag{3}$$

Here the coupling constant G_3 has dimension (-2) and is related to an UV cut-off Λ, reflecting the scale of chiral symmetry breaking of the order of 1 GeV. Note that it is the heavy quark residual momentum $(P_Q - m_Q\,v^\mu)^2 \sim O(p_q^2) < \Lambda^2$ which is regularized, and in this way the NJL model may indeed be applied. Note also, that all reference to gluons is gone, and consequently the model has no idea about confinement.

The above lagrangian can be transformed *exactly* into a quark-meson interaction by integrating in auxiliary fields, which in the HQL leads to

$$\tilde{\mathcal{L}}^{\bar{\chi}Q} \xrightarrow{m_Q\to\infty} -\bar{Q}_v\,[H+K]\,\chi - \bar{\chi}\,\left[\bar{H}+\bar{K}\right]\,Q_v + \frac{1}{2G_3}\,\mathrm{Tr}\left[\bar{H}H - \bar{K}K\right]\ . \tag{4}$$

Here we introduced the above mentioned non-linear representation by rotating the light quark fields $\chi_R = \xi\,q_R\ ,\ \ \chi_L = \xi^\dagger\,q_L$. The Dirac matrices for the spin-symmetry doublets with $J^P = (0^-, 1^-)$ and $J^P = (0^+, 1^+)$, respectively, are given by

$$H \equiv \frac{1+\not{v}}{2}\,(i\Phi_5\gamma_5 + \not{\Phi})\ , \qquad K \equiv \frac{1+\not{v}}{2}\,(\Phi + i\not{\Phi}'_5\gamma_5)\ , \tag{5}$$

with $v_\mu\,\Phi^\mu = v_\mu\,\Phi^\mu_5 = 0$. The quark fields are now integrated out explicitly by a functional Gaussian integration. The real part of the resulting quark determinant after Wick-rotation $(D \to D_E)$ is regularized by a proper-time integral

$$\ln|\det D| \ \to\ -\frac{N_c}{2}\int_{1/\Lambda^2}^{1/\mu^2}\frac{ds}{s}\int \mathrm{Tr}\left[e^{-s\,D_E^\dagger D_E}\right]\ . \tag{6}$$

Here, μ serves as an adjustable parameter and may be interpreted as the scale up to which the quarks have been integrated out. Especially, at $\mu = \Lambda$ the contribution of the quark determinant vanishes by construction, and we recover the original interaction of quarks with static meson fields in Eq. (4).

It is a straight forward task to generalize the presented ideas to heavy mesons of higher excitations like the $J^P = (1^+, 2^+)$ multiplet. It can be represented as[5]

$$T^\mu = \frac{1+\not{v}}{2}\left\{\Phi^{\mu\nu}\gamma_\nu + \sqrt{\frac{3}{2}}\Phi_5{}^\nu\, i\gamma_5\,\left(g_{\mu\nu} - \frac{1}{3}\gamma_\nu(\gamma_\mu - v_\mu)\right)\right\} \tag{7}$$

with $v_\nu\Phi^{\mu\nu} = v_\nu\Phi_5{}^\nu = 0$, $\Phi^{\mu\nu} = \Phi^{\nu\mu}$ and $\Phi^\mu_\mu = 0$. For the light degrees of freedom with $j_l = 3/2$ we construct a Rarita-Schwinger representation from the light quark fields, introducing a covariant derivative $i\mathcal{D}_\nu \equiv i\partial_\nu + i/2\,(\xi^\dagger\partial_\nu\xi + \xi\partial_\nu\xi^\dagger)$ of the chiral $SU(3)_V$,

$$\tilde{\mathcal{L}}^{\chi Q_v} \quad \to \quad -\bar{Q}_v\, T^\nu\, i\overrightarrow{\mathcal{D}}_\nu\, \chi + \bar{\chi}\, i\overleftarrow{\mathcal{D}}_\mu\, \bar{T}^\mu\, Q_v - \frac{1}{2G_4}\,\mathrm{Tr}\left[\bar{T}^\mu T_\mu\right] \;, \tag{8}$$

introducing a new coupling constant $G_4 \sim O(1\ \mathrm{GeV})^{-4}$. For further details we refer to our original works[6,7].

3. Results



Figure 1: The solutions m_q of Eq. (9) as an implicit function of the scale μ.

For low μ the contribution of the quark determinant leads to dynamical breaking of chiral symmetry. In the NJL model this is governed by the gap-equation for the light quark constituent mass[8]

$$m_q = m_q^{(0)} + 8\,G_1\, m_q\, I_1 \;, \qquad I_1 = \frac{N_c}{16\,\pi^2}\left(\Lambda^2 - \mu^2 + O(m_q^2)\right) \;. \tag{9}$$

Fig. 1 shows the behaviour of the solution for m_q as a function of the scale μ in the chiral limit, i.e. for a vanishing current quark mass $m_q^{(0)} = 0$. We may use μ for separating the IR region of the proper-time integral in Eq. (6) – which is believed to be governed by confinement and can not be well described in our model – from the region where the model is assumed to lead to reasonable results, namely to a consistent description of dynamical chiral symmetry breaking. The dashed vertical line indicates the convenient choice $\mu = 300$ MeV. Note that in the chiral limit the critical scale for the phase transition can be read off analytically from Eq. (9)

$$\mu_c^2 \;=\; \Lambda^2 - \frac{2\,\pi^2}{3\,G_1} \approx (550\ \mathrm{MeV})^2 \;, \qquad (G_1 = 5.25\ \mathrm{GeV}^{-2},\ \Lambda = 1.25\ \mathrm{GeV})^6 \;. \tag{10}$$

With the self-energy contributions of the quark-determinant for scales $\mu < \Lambda$ the several meson fields in the NJL model become dynamical degrees of freedom. We stress, that the limit $\mu \to 0$ in the proper-time integrals is in general not suitable if the external momenta exceed the sum of the internal quark masses since by analytic continuation the self-energy then receives an imaginary part, which by the optical theorem is connected with unphysical quark-antiquark thresholds.

The mass-spectrum of the heavy mesons in the NJL model is then determined by the value of the 4-quark coupling constant G_3 (with all other parameters fixed in the light meson sector). The NJL model further gives simple predictions for the weak decay constants of the heavy pseudoscalar (P) and vector mesons (V) in terms of the wave function renormalization factors‡ Z and the coupling constant G_3,

$$\left.\begin{array}{l}\sqrt{M_P}\, f_P = (1+\delta)\,\dfrac{\sqrt{Z_P/M_P}}{G_3}\\[2ex] \sqrt{M_V}\, f_V = \dfrac{\sqrt{Z_V/M_V}}{G_3}\end{array}\right\} \xrightarrow{\text{HQL}} \sqrt{M_H}\, f_H = \frac{\sqrt{Z_H}}{G_3} \tag{11}$$

Here, δ denotes the contribution from the mixing of the pseudoscalar and axial-vector states beyond the HQL. The perturbative corrections to Eq. (11) are known; in the leading-log approximation one has[9] $f_D = f_B\ (\alpha_s(m_c)/\alpha_s(m_b))^{6/25}$.

The results are collected in Table 1. Compared to experiment the $SU(3)_F$-mass splitting due to the different light quark masses $m_u = 300$ MeV and $m_s = 510$ MeV is somewhat underestimated. The experimental fine-splitting between the spin-symmetry partners is recovered for rather small values of m_b and m_c. We further observe large corrections to the HQL scaling law for the pseudoscalar decay constant, whereas the deviations for the vector decay constant are within the usual expectation. This is mostly due to the large value of the mixing contribution δ which obtains values up to 50% in the case of D-mesons. The values for the weak decay constants are on the lower side of the range obtained from lattice results[10].

Table 1: Heavy Meson masses and decay constants in the NJL model for a) the HQL and b) finite values of m_Q. Here $\bar\Lambda$ is the mass-difference between heavy meson and heavy quark in the HQL. The heavy quark masses are fitted to the averaged heavy meson masses $\bar M_H = 1/4\,(3\,M_V + M_P)$. These are then used to estimate the mass-splittings $\delta_H = M_V - M_P$ and the decay constants.

a)

G_3	$\bar\Lambda_u$	$\bar\Lambda_s - \bar\Lambda_u$	$M_K - M_H$	f_B	f_{B_s}/f_B
$[\text{GeV}]^{-2}$		[MeV]		[MeV]	
6.3	300	60	245	140	1.04
4.5	400	75	190	150	1.04
2.9	500	80	140	160	1.03
Exp.[12]		100			

b)

G_3	m_b	m_c	δ_B	δ_D	f_B	f_{B^*}	f_D	f_{D^*}
$[\text{GeV}]^{-2}$	[GeV]		[MeV]			[MeV]		
6.3	4.97	1.53	81	239	130	135	160	200
4.5	4.88	1.43	63	192	140	145	170	225
2.9	4.78	1.33	46	146	150	155	190	260
Exp.[12]			46	141				

The NJL result for the Isgur-Wise function is obtained from the quark determinant by insertion of a heavy quark current. We present the NJL results for the slope parameter

---

‡Note that in the HQL a factor of M_H is absorbed into the normalization of the heavy meson fields.

$\rho = \sqrt{-\xi'(1)}$ and the curvature $c_0 = \xi''(1)/2$,

$$\begin{aligned} v\cdot p = 300 \text{ MeV}: &\quad \rho = 0.84\ , \quad c_0 = 0.49\ ; \\ v\cdot p = 400 \text{ MeV}: &\quad \rho = 0.94\ , \quad c_0 = 0.72\ ; \\ v\cdot p = 500 \text{ MeV}: &\quad \rho = 1.07\ , \quad c_0 = 1.18\ . \end{aligned} \tag{12}$$

Note that here the values of $v{\cdot}p = \bar\Lambda$ and ρ and c_0 are positively correlated. This should be compared with the estimated theoretical relation[11] $c_0 \simeq 0.72\,\rho^2 - 0.09$ together with the phenomenological bounds $0.84 \le \rho \le 1.00$.

4. Conclusions

We have presented the NJL quark model as a low-energy approximation to QCD including the approximate symmetries for light and heavy flavors. It has been shown that this model is equivalent to a relativistic quark-meson model for the several heavy meson states. We have introduced an adjustable scale parameter μ in the proper-time regularized quark determinant, such that at $\mu = \Lambda$ the model matches to the case of static meson fields which become then dynamical for $\mu < \Lambda$. The dynamical breaking of chiral symmetry occurs around $\mu \approx 550$ MeV. We then chose a finite value of $\mu \approx 300$ MeV in order to separate the IR region of the proper-time integration. With this one obtains reasonable results for the heavy meson mass spectrum both, with finite and infinite heavy quark masses, for the decay constants and for the Isgur-Wise form factors.

Acknowledgements

I would like to express my gratitude to the Organizing Committee of the workshop and the people at the JINR Dubna for their warm hospitality as well as to the Heisenberg-Landau Program for financial support. It is also a pleasure to thank D. Ebert, H. Reinhardt and R. Friedrich for their contributions to the results presented here.

References

1. see e.g. T. Mannel, W. Roberts and Z. Ryzak, *Nucl. Phys.* **B368** (1992) 204.
2. A. F. Falk, *Nucl. Phys.* **B378** (1992) 79.
3. G. Burdman and J. F. Donoghue, *Phys. Lett.* **B280** (1992) 287;
   M. B. Wise, *Phys. Rev.* **D45** (1992) R2188.
4. R. Casalbuoni et al., hep-ph/9605342 (1996).
5. J. G. Körner, D. Pirjol and K. Schilcher, *Phys. Rev.* **D47** (1993) 3955;
   P. Cho and S. P. Trivedi, *Phys. Rev.* **D50** (1994) 381.
6. D. Ebert, T. Feldmann, R. Friedrich and H. Reinhardt, *Nucl. Phys.* **B434** (1995) 619.
7. D. Ebert, T. Feldmann and H. Reinhardt, in preparation.
8. D. Ebert and H. Reinhardt, *Nucl. Phys.* **B271** (1986) 188.
9. see e.g. H. Politzer and M. B. Wise, *Phys. Lett.* **B206** (1988) 681.
10. H. Wittig, contribution to this workshop, hep-ph/9606371, and refs. therein.
11. M. Neubert, hep-ph/9511409; I. Caprini and M. Neubert, hep-ph/9603414.
12. L. Montanet et al. (Particle Data Group), *Phys. Rev.* **D50** (1994), 1173.